



ELSEVIER

Contents lists available at ScienceDirect

Zoologischer Anzeiger

journal homepage: www.elsevier.com/locate/jcz

Short communication

A new record of the marine chaetonotid gastrotrich genus *Diuronotus* (Paucitubulatina: Muselliferidae) from the Sea of JapanNiels W.L. Van Steenkiste ^{a, b, *}, Kevin C. Wakeman ^{c, d}, Brian S. Leander ^{a, b}^a Department of Botany, University of British Columbia, 3156-6270 University Blvd, Vancouver, BC, V6T 1Z4, Canada^b Department of Zoology, University of British Columbia, 4200-6270 University Blvd, Vancouver, BC, V6T 1Z4, Canada^c Institute for the Advancement of Higher Education, Hokkaido University, North 17, West 8, Kita-ku, Sapporo, 060-0808, Japan^d Graduate School of Science, Hokkaido University, North 10, West 8, Kita-ku, Sapporo, 060-0810, Japan

ARTICLE INFO

Article history:

Received 27 November 2020

Received in revised form

21 December 2020

Accepted 22 December 2020

Available online 24 December 2020

Keywords:

Chaetonotida

Gastrotricha

Hokkaido

Species distribution

Sea of Japan

ABSTRACT

Marine gastrotrichs of the Pacific Ocean are poorly known. Here, we report on the finding of a marine chaetonotid gastrotrich of the genus *Diuronotus* from an intertidal beach within the Sea of Japan in Hokkaido (Japan). The Japanese individual shows a very close resemblance to *Diuronotus aspetos*. This new record is a consequential extension of its biogeographic range; previous records for representatives of this genus are confined to West Greenland, the North Sea and the east coast of North America. This rarely encountered, but seemingly widespread genus of marine gastrotrichs exemplifies our limited understanding of meiofaunal diversity and distribution patterns caused by sampling bias and insufficient knowledge on nominal species complexes.

© 2020 Elsevier GmbH. All rights reserved.

1. Introduction

The only two described species of the gastrotrich genus *Diuronotus* Todaro, Balsamo & Kristensen, 2005, *Diuronotus aspetos* Todaro, Balsamo & Kristensen, 2005 and *Diuronotus rupperti* Todaro, Balsamo & Kristensen, 2005, are relatively large marine members of the Paucitubulatina that can easily be recognized by their diagnostic posterior furca with one pair of primary and one pair of secondary adhesive tubes. Their cuticle is ornamented with heteromorphic keeled scales with a concave posterior edge. *D. aspetos* was originally found in shallow subtidal sand under the mid-winter sea ice around Disko Island, Greenland (Todaro et al. 2005). Subsequently, *D. aspetos* has been reported three more times — twice from a similar habitat in the same area of Greenland during the Arctic summer (Balsamo et al. 2010; Bekhouche & Vorsaae 2016), and once from subtidal sand in the temperate southeastern part of the North Sea (Kieneke 2015) (Fig. 1A). *D. rupperti* was collected from intertidal sand in the Kattegat between the North Sea and the Baltic Sea, while a third, hitherto unidentified species originating from the east coast of North

America, is also attributed to the genus *Diuronotus* (Todaro et al. 2005) (Fig. 1A).

We report on the finding of one specimen of *Diuronotus*, which closely resembles *D. aspetos*, from Hokkaido, Japan, along the northeastern edge of the Sea of Japan (Fig. 1A).

2. Material and methods

One individual of *Diuronotus* was collected in February 2019 from the lower intertidal zone of a sandy beach in Otaru, Hokkaido, Japan (43°13'33"N, 141°00'58"E). Sand from the top layer was collected during low tide and transferred to the lab at Hokkaido University. Live meiofauna was separated from the sand using the MgCl₂ anesthetization-decantation method (Todaro et al. 2019). The individual was isolated and observed under a stereoscope and subsequently whole mounted alive to be studied, photographed and measured under an Olympus BX53 microscope equipped with DIC optics. Micrographs were edited with Adobe Photoshop v21.1 for contrast and brightness and the figure was put together in Adobe Illustrator v24.1.

* Corresponding author. Department of Botany, University of British Columbia, 3156-6270 University Blvd, Vancouver, BC, V6T 1Z4, Canada.

E-mail addresses: niels_van_steenkiste@hotmail.com, niels.vansteenkiste@botany.ubc.ca (N.W.L. Van Steenkiste).




Fig. 1. *Diuronotus* cf. *aspetus* from the Sea of Japan (A) Global distribution of all representatives of the genus *Diuronotus* including the new record from the Sea of Japan (B–C) Live specimen slightly squeezed under cover slip in two different focal planes (ventral and central) (D) Anterior end (E) Detail of keeled cuticular scales covering the body. Abbreviations: in, intestine; ph, pharynx; sTbP, secondary adhesive tube; TbP, primary adhesive tube. White arrows indicate cuticular rods protruding through the mouth ring. Scale bars in DIC micrographs: B–C = 100 µm; D–E = 20 µm.

The specimen was preserved in lactophenol as a permanent whole mount, which was deposited in the Hokkaido University Museum, Japan (accession number HokkNVS 2019).

3. Results and discussion

The overall morphology is very similar to that of *D. aspetos*. Measurements of the Japanese specimen are as follows: body length 580 µm, pharynx (ph) length 152 µm, fleshy basal portions of the furcal branches 62 µm, primary adhesive tubes (TbP) 23 µm, secondary adhesive tubes (sTbP) 60 µm, scales up to 15 µm long (18 µm including the spine of the keel) and 5 µm wide. Pharyngeointestinal junction at U28.7. These values are similar or slightly larger than those from the adult specimens of *D. aspetos* from Greenland, and larger than those from the smaller subadult specimen also attributed to *D. aspetos* from the North Sea. External diagnostic features including the keeled scales with posterior concave edge and the four posterior adhesive tubes — the two secondary tubes three times as long as the two primary tubes — are clearly visible (Fig. 1B–E). The relative and absolute length of the

secondary tubes clearly distinguishes our specimen from *D. rupperti*, for which these tubes are up to 8.5 µm in adults and completely lacking in juveniles (Todaro et al. 2005). Apart from the outline of the pharynx and intestine, no additional features of the internal organs were observed, possibly because the individual from Japan is subadult. The internal morphology and phylogenetic position of *D. aspetos* have been extensively studied in Greenland specimens (Balsamo et al. 2010; Bekkouche & Worsaae 2016; Todaro et al. 2005).

A total of 30 species of gastrotrichs have been reported from the Sea of Japan as defined by the International Hydrographic Organization (IHO) (Table 1). Almost all these records come from the extreme southeastern edge of the Sea of Japan (east coast of Korea) and its transition zone to the Yellow Sea and East China Sea (south coast of Korea, Jeju Island and the Korea Strait) (e.g. Lee et al. 2017 and references therein). However, this warmer transition zone is not included in the Sea of Japan when it is defined as a marine ecoregion (Spalding et al. 2007). When considering the latter definition of the Sea of Japan, the number of gastrotrich species is reduced to 15 (Table 1). Only three other species of gastrotrichs

Table 1

Records and distribution of marine gastrotrichs in the Sea of Japan as defined by the International Hydrographic Organization (IHO) or defined as a marine ecoregion (ME) (Spalding et al. 2007). Abbreviations: EC, East Coast; HK, Hokkaido; JI, Jeju Island; SC, South Coast; TI, Tsushima Island; PG, Peter the Great Bay.

Taxa	Sea of Japan (IHO)	Sea of Japan (ME)
Macrodasyida		
<i>Acanthodasys comptus</i> Lee, 2012	Korea, JI	—
<i>Cephalodasys mahoae</i> Yamauchi & Kajihara, 2018	Japan: HK	Japan: HK
<i>Crasiella clauseni</i> Lee & Chang, 2012	Korea: JI	—
<i>Dendrodasys duplex</i> Lee, Chang & Kim, 2014	Korea: EC, JI	Korea: EC
<i>Diplodasys ankelii</i> Wilke, 1954	Korea: JI	—
<i>Diplodasys meloriae</i> Todaro, Balsamo & Tongiorgi, 1992	Korea: EC, JI	Korea: EC
<i>Lepidodasys laeviacus</i> Lee & Chang, 2011	Japan: TI	—
<i>Lepidodasys tsushimaensis</i> Lee & Chang, 2011	Japan: TI	—
<i>Pseudostomella longifurca</i> Lee & Chang, 2002	Korea: EC, JI	Korea: EC
<i>Ptychostomella jejuensis</i> Lee, Hwang & Chang, 2009	Korea: JI	—
<i>Ptychostomella orientalis</i> Lee & Chang, 2003	Korea: EC	Korea: EC
<i>Ptychostomella papillata</i> Lee & Chang, 2003	Korea: EC	Korea: EC
<i>Tetranchyroderma aethesbregmum</i> Lee & Chang, 2012	Korea: JI	—
<i>Tetranchyroderma coreense</i> Lee, Kim & Chang, 2013	Korea: JI	—
<i>Tetranchyroderma gracilium</i> Chang, Lee & Clausen, 1998	Korea: JI	—
<i>Tetranchyroderma heterotentaculatum</i> Chang & Lee, 2001	Korea: EC, JI	Korea: EC
<i>Tetranchyroderma hoonsooi</i> Chang & Lee, 2001	Korea: EC, JI, SC	Korea: EC
<i>Tetranchyroderma hummoni</i> Lee, Chang & Kim, 2017	Korea: EC	Korea: EC
<i>Tetranchyroderma insolitum</i> Lee & Chang, 2012	Korea: SC	—
<i>Tetranchyroderma monokerosum</i> Lee & Chang, 2007	Korea: JI	—
<i>Tetranchyroderma multicirratum</i> Lee & Chang, 2007	Korea: JI	—
<i>Tetranchyroderma oblongum</i> Lee, Kim & Chang, 2013	Korea: EC	Korea: EC
<i>Tetranchyroderma pinnatum</i> Lee & Chang, 2014	Korea: SC	—
<i>Tetranchyroderma schizocirratum</i> Chang, Kubota & Shirayama, 2002	Korea: JI	—
<i>Thaumastoderma appendiculatum</i> Chang, Lee & Clausen, 1998	Korea: EC	Korea: EC
<i>Thaumastoderma copiophorum</i> Chang, Lee & Clausen, 1998	Korea: JI	—
<i>Turbanella lobata</i> Yamauchi & Kajihara, 2018	Japan: HK	Japan: HK
Chaetonotida		
<i>Aspidiophorus oculatus</i> Todaro et al., 2009	Russia: PG	Russia: PG
Diuronotus cf. aspetos	Japan: HK	Japan: HK
<i>Halichaetonotus aculifer</i> (Gerlach 1953)	Korea: EC, JI	Korea: EC
<i>Halichaetonotus atlanticus</i> Kisielewski, 1988	Korea: EC, JI, SC	Korea: EC

have been reported from the northern part of the Sea of Japan: *Aspidiophorus oculatus* Todaro et al. 2009 from Peter the Great Bay, Russia, and *Cephalodasys mahoae* Yamauchi & Kajihara, 2018 and *Turbanella lobata* Yamauchi & Kajihara, 2018 from Hokkaido, Japan (Todaro et al. 2009; Yamauchi & Kajihara 2018). This makes *Diuronotus* cf. *aspertos* the third marine gastrotrich and first chaetonotid found along Hokkaido's coastline on the Sea of Japan. For other coastal areas of Japan (Pacific Ocean, Korea Strait, Seto Inland Sea), twelve species of gastrotrichs have been reported (Yamauchi & Kajihara 2018 and references therein).

Upon its discovery under the sea ice in Greenland, *D. aspetos* was considered a high Arctic species. However, the additional record from the North Sea attributed to this species indicated a much wider distribution and a wider temperature tolerance for this species (Kieneke 2015). Our finding of *D. cf. aspetos* in the Sea of Japan suggests that this species might not only occur in the Arctic and temperate Northern Atlantic, but also in the temperate Northern Pacific. However, increased collection efforts in the North Sea, the Sea of Japan and other seas of the northern hemisphere and a more comprehensive integrative taxonomic approach are imperative to test whether the singletons from Germany and Japan indeed belong to *D. aspetos* or are closely related sister species and, in case of the first scenario, whether its Arctic and circumtemperate distribution is either disjunct or continuous. Current data on the diversity and distribution of gastrotrich species recognize cosmopolitan and trans-oceanic distributions for several species, but also high levels of endemism in other species (Garraffoni & Balsamo 2017). Sampling bias — especially in the Pacific Ocean — and insufficient knowledge on the existence of cryptic species complexes in widespread nominal species need to be addressed to better understand species diversity and global distribution patterns of marine gastrotrichs.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

NWLV and BSL were funded by the Hakai Institute and the National Sciences and Engineering Research Council of Canada (NSERC 2019-03986). KCW was supported by the Japanese Society for the Promotion of Science (18K14774).

References

- Balsamo, M., Guidi, L., Ferraguti, M., Pierboni, L., Kristensen, R.M., 2010. *Diuronotus aspetos* (Gastrotricha): new morphological data and description of the spermatozoon. *Helgol. Mar. Res.* 64, 27–34. <https://doi.org/10.1007/s10152-009-0163-x>.
- Bekkouche, N., Worsaae, K., 2016. Neuromuscular study of early branching *Diuronotus aspetos* (Paucitubulatina) yields insights into the evolution of organs systems in Gastrotricha. *Zool. Lett.* 2, 21. <https://doi.org/10.1186/s40851-016-0054-3>.
- Garraffoni, A.R.S., Balsamo, M., 2017. Is the ubiquitous distribution real for marine gastrotrichs? Detection of areas of endemism using Parsimony Analysis of Endemicity (PAE). *Proc. Biol. Soc. Wash.* 130, 198–211. <https://doi.org/10.2988/17-00011>.
- Kieneke, A., 2015. Record of the 'Arctic' marine gastrotrich *Diuronotus aspetos* (Paucitubulatina) from the southern North Sea. *Mar. Biodivers.* 45, 615–616. <https://doi.org/10.1007/s12526-014-0294-4>.
- Lee, J., Chang, C., Kim, I.-H., 2017. A new species of the genus *Tetranchyroderma* (Gastrotricha: macrodasyida: Thaumastodermatidae) from Korea, with a peculiar pentangular armature. *Proc. Biol. Soc. Wash.* 130, 156–165. <https://doi.org/10.2988/17-00016>.
- Spalding, M.D., Fox, H.E., Allen, G.R., Davidson, N., Ferdaña, Z.A., Finlayson, M., Halpern, B.S., Jorge, M.A., Lombana, A., Lourie, S.A., Martin, K.D., McManus, E., Molnar, J., Recchia, C.A., Robertson, J., 2007. Marine ecoregions of the world: a

Bioregionalization of coastal and shelf areas. *Bioscience* 57, 573–583. <https://doi.org/10.1641/B570707>.

Todaro, M.A., Balsamo, M., Kristensen, R.M., 2005. A new genus of marine chaetodontids (Gastropoda), with a description of two new species from Greenland and Denmark. *J. Mar. Biol. Assoc. U. K.* 85, 1391–1400. <https://doi.org/10.1017/S0025315405012579>.

Todaro, M.A., Dal Zotto, M., Maiorova, A.S., Adrianov, A.V., 2009. A new species of *Aspidiophorus* (Gastropoda, Chaetodontida) from the Russian Far East with a key to marine species of the genus. *Mar. Biol. Res.* 5, 297–303. <https://doi.org/10.1080/17451000802419430>.

Todaro, M.A., Sibaja-Cordero, J.A., Segura-Bermúdez, O.A., Coto-Delgado, G., Goebel-Otárola, N., Barquero, J.D., Cullell-Delgado, M., Dal Zotto, M., 2019. An introduction to the study of Gastropoda, with a taxonomic key to families and genera of the group. *Diversity* 11, 117. <https://doi.org/10.3390/d11070117>.

Yamauchi, S., Kajihara, H., 2018. Marine macrodasyida (Gastropoda) from Hokkaido, northern Japan. *Species Divers.* 23, 183–192. <https://doi.org/10.12782/specdiv.23.183>.